


ADULT PRETRIAL TEST  
\*\*\*\*\*

NAME: Example Report  
ID# : 0000000000  
AGE : 31 SEX: Male  
ETHNICITY/RACE: Caucasian  
EDUCATION/HIGHEST GRADE: 12  
MARITAL STATUS: Married  
DATE OF BIRTH: 01/01/1979  
DATE SCORED: 12/11/2010

CONFIDENTIAL REPORT

Adult Pretrial Test (APT) results are working hypotheses and are to be used in conjunction with an interview, record review and experienced staff judgment. No diagnosis or recommendation should be based solely upon APT results.



\*\*\* Substance (Dependency/Abuse) Classification: substance abuse

ADDITIONAL INFORMATION PROVIDED BY CLIENT

Age at first arrest.....	18	Total number of times arrested...	2
Age at first conviction.....	18	Jail sentences.....	1
Felony arrests.....	1	Prison sentences.....	1
Times on probation.....	1	Alcohol-related arrests.....	1
Probation revocations.....	0	Drug-related arrests.....	1
Times on parole.....	0	DUI or DWI arrests.....	1
Parole revocations.....	0	Months employed in last 12 months	12

## \* \* SUMMARY PARAGRAPHS EXPLAINING CLIENT'S ATTAINED SCALE SCORES \* \*

## TRUTHFULNESS SCALE: MEDIUM RISK RANGE

RISK PERCENTILE:55

This APT profile is accurate. However, there is a tendency for this defendant to deny common problems and to portray self in an overly favorable light. Specific questions will usually be answered more accurately than open-ended or general type questions. This is an accurate APT profile and other APT scale scores are accurate. This defendant has adequate reading skills. Responses are reasonably accurate.

DEPENDENCY/ABUSE CLASSIFICATION: Three or more of the seven DSM-IV dependency items (or their equivalents) are not admitted to, consequently the classification of 'dependency' as defined here does not apply. Moreover, this person reports (item #74) that he or she has not been diagnosed 'substance dependent' in the past. One or more of the substance abuse items (or their equivalents) are admitted to, which meets the criteria of 'substance abuse' as defined here. These admissions include: 140, 143, 59. This client describes their drinking as a moderate problem and their drug use as a minimum problem. This person's Alcohol Severity Scale score is elevated (at or above the 70th percentile) whereas their Drugs Severity Scale score is not (at or below the 69th percentile) elevated. This APT profile is not uncommon in the early stages of substance use, or after relapse. In summary, this client meets the DSM-IV substance abuse classification. And within this abuse category their degree of risk is represented by their Alcohol Severity Scale score.

## ALCOHOL SEVERITY SCALE: PROBLEM RISK RANGE

RISK PERCENTILE:87

This defendant's Alcohol Severity Scale score is in the Problem Risk (55 to 89th percentile) range. Problematic alcohol-related disorders are not identified by chance. This level of severity warrants consideration of outpatient treatment (individual or group counseling) services. The duration for outpatient treatment will be to a large extent contingent upon defendant compliance and responsiveness to treatment. Mandatory treatment attendance is recommended on a weekly basis for a period of six months. Should the primary care psychiatrist, psychologist or counselor decide to terminate treatment earlier than six months, they would notify the court in writing. Should the defendant want to continue treatment beyond six months, it would be his/her responsibility to arrange for said services. Family counseling, Alcoholics Anonymous (AA) and Rational Recovery (RR) meetings might augment treatment but should not take the place of outpatient treatment. Either this defendant is a "recovering" alcoholic or this person meets the criteria for an alcohol or drinking problem. With regard to alcohol, this defendant meets the problem risk criteria. Although this person does not meet the DSM-IV dependency criteria, their scale score indicates that the person does represent a substance use risk.

## DRUGS SEVERITY SCALE: PROBLEM RISK RANGE

RISK PERCENTILE:63

The defendant is in the early stages of an illicit drug-related disorder, as their Drugs Severity Scale score is in the Problem Risk (55 to 89th percentile) range. The appropriate level of treatment for this problem severity is outpatient (individual counseling, group counseling or outpatient addiction treatment) services on a minimum once-a-week schedule. Length of treatment will vary contingent upon

defendant compliance and response to treatment. Mandatory outpatient treatment is recommended for six months. The defendant could, if desired, continue treatment on his/her own beyond six months. Initiation of treatment and completion of treatment reports (including diagnosis, dates of attendance and prognosis) should be submitted to the court designee. Family counseling or involvement in the defendant's treatment plan should be encouraged. Narcotics Anonymous (NA) and Cocaine Anonymous (CA) meetings might augment--but not take the place of--outpatient addiction services. This defendant's pattern of Drugs Severity Scale responses meets the criteria of problem severity or risk. With regard to drugs, this defendant meets the problem risk criteria. Although this person does not meet the DSM-IV dependency criteria, their scale score indicates that the person does represent a substance use risk.

ANTISOCIAL REACTION SCALE: PROBLEM RISK RANGE RISK PERCENTILE:89

An established pattern of antisocial thinking and attitude is evident. Problem risk is characterized by many antisocial attitudes and behaviors as well as difficulty maintaining responsible relationships and loyalties. These individuals are frequently callous, irresponsible, and lack a foundation of mutual affection or trust. Many are boastful, deceitful and given to tantrums or outbursts of rage. Poor work histories, nonpayment of bills and difficulty conforming to social norms are common. With regard to the Antisocial Reaction Scale, this defendant attained a problem risk score.

LETHALITY SCALE: MAXIMUM RISK RANGE RISK PERCENTILE:93

Violent tendencies are indicated and a pattern of violence is likely well established. This person could be dangerous to self or others. Violent behavior is characterized by ruthlessness, savageness, destructiveness and explosiveness. Substance abuse, jealousy and perceived stress could escalate into violent behavior. This defendant would likely be intimidating, threatening and potentially dangerous. This is a violent person. Prior violence is likely. This defendant would benefit from anger management group counseling. And, consideration might be given to obtaining a more comprehensive psychological evaluation (diagnosis, treatment plan and prognosis).

STRESS QUOTIENT SCALE: MEDIUM RISK RANGE RISK PERCENTILE:65

Stress coping abilities are not well established, however, stress does not present as a focal issue. Stress-related problems are characterized by irritability, instability and interpersonal conflict. Coping with stress is a process of adaptation. Symptoms of stress can be psychological or physiological and include anxiety, depression, irritability, substance abuse and moodiness. This is a medium risk stress coping score.

APT RESPONSES

-----

1- 50	FTTFFTTFFT	FFTTFFTTFF	TTFETTTFFT	FETTTFFFTT	FETTFTTFFT
51-100	TFTTFFTTTF	FFTTFFTTFT	TFTTFFTTTF	TFTTFFTTFT	TFTTFFTTFT
101-150	FTTTT22331	2122321312	2313223134	1232423222	4434444444
151-162	4423444441	24			

SIGNIFICANT ITEMS: These answers are the defendant's self-reported responses. And, they represent direct admissions or unusual responses, which may help in understanding the defendant's situation.

ALCOHOL SEVERITY

-----

- 2. States concerned about drinking
- 34. Reports a recovering alcoholic
- 44. Drinking problem in last year
- 55. States serious drinking problem
- 70. Must overcome drinking problem
- 81. Asked for help with drinking
- 87. Has been treated for drinking

DRUGS SEVERITY

-----

- 13. Admits smokes pot to relax
- 30. Has attended NA or CA meetings
- 51. Has sought drug problem help
- 83. Presently in drug treatment
- 100. Admits is dependent on drugs

ANTISOCIAL REACTION

-----

- 29. Does not trust many people
- 40. People tell client: antisocial
- 50. Difficult keeping relationships
- 63. Gets in trouble with law a lot
- 90. People think client antisocial
- 96. Lot of needless lying/conning

LETHALITY (VIOLENCE)

-----

- 5. Client denies being nonviolent
- 26. More dangerous than age peers
- 37. Can be dangerous when angered
- 73. Arrest: assault/violent crime
- 79. Has deliberately hurt someone
- 103. Often thinks of death/suicide

STRUCTURED INTERVIEW (DEFENDANT'S SELF-REPORT):

-----

- 140. Repeated substance use: neglecting children or responsibilities
- 141. Did not use substances before driving or in dangerous settings
- 142. Denies repeated substance use resulted in legal problems
- 143. Continued substance use despite family arguments & social problems
- 144. Denies serious substance problems or knowing it caused problems
- 145. States has not used more to get intoxicated or high
- 146. Denies withdrawal symptoms after reducing/stopping substance use
- 147. Did not use alcohol or drugs to avoid/reduce withdrawal symptoms
- 148. Denies using alcohol or drugs more or longer than intended
- 149. Has not tried and failed to control or stop using alcohol/drugs
- 150. Denies time is spent obtaining/using/recovering from alcohol/drugs
- 151. Not high, drunk or in withdrawal before/during school/work/driving
- 152. Denies reducing activities or family because of alcohol or drugs
- 153. Describes drinking as a moderate problem
- 154. Client states little motivation for alcohol treatment or help
- 155. States has not been enrolled in an alcohol treatment program
- 156. Reports is not a "recovering" person
- 157. States drug use is not a problem
- 158. Has not been enrolled in a drug treatment program
- 159. Client states has no need for drug treatment or help
- 160. Client reports has serious family problems
- 161. Client reports arrest for crime involving force or threat of force
- 162. Has not been suicidal/homicidal in last 6 months

RECOMMENDATIONS: \_\_\_\_\_

\_\_\_\_\_

STAFF MEMBER SIGNATURE

DATE

(APT TEST # 1 )